arXiv:1602.00316v1 [math.CA] 31 Jan 2016

# On Certain Generalizations of Rogers-Ramanujan Type Identities

Ahmad El-Guindy* Mourad E.H. Ismail†

**Abstract**

We state and prove a number of unilateral and bilateral $q$-series identities and explore some of their consequences. Those include certain generalizations of the $q$-binomial sum which also generalize the $q$-Airy function introduced by Ramanujan, as well as certain identities with an interesting variable-parameter symmetry based on limiting cases of Heine's transformation of basic hypergeomteric functions.

MSC (2010): Primary 33D15 ; Secondary 33D70
Keywords: Basic Hypergeometric Functions, Heine Transform, Ramanujan's $q$-Airy Function, Bilateral $q$-series

## 1 Introduction

The theory of $q$-series is well known for a number of fascinating identities with far reaching number theoretic consequences. A famous such example comes from the Rogers-Ramanujan identities

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q;q)_n} = \frac{1}{(q,q^4;q^5)_\infty} \tag{1.1}$$

$$\sum_{n=0}^{\infty} \frac{q^{n^2+n}}{(q;q)_n} = \frac{1}{(q^2,q^3;q^5)_\infty}, \tag{1.2}$$

where we follow the standard notations for $q$-shifted factorials and basic hypergeometric series as in the books [4], [6], [7]. References for the Rogers-Ramanujan identities, their origins and many of their applications are in [1], [2], and [4]. In particular we recall the partition theoretic interpretation of the first Rogers–Ramanujan identity as the partitions of an integer $n$ into parts $\equiv 1$ or $4 \pmod 5$ are equinumerous with the partitions

*Research supported by a QNRF grant from Qatar Foundation NPRP No. : 7-1360-1-254
†Research supported by a QNRF grant from Qatar Foundation NPRP No. : 7-1360-1-254

of $n$ into parts where any two parts differ by at least 2. The Roger-Ramanujan identities had many extensions and generalizations to different settings. One noteworthy generalization is to extend the identities in (1.2) to evaluate the sum $\sum_{n=0}^{\infty} q^{n^2+mn}/(q;q)_n$, $m = 0, \pm 1, \pm 2, \cdots$, see [5]. One can view the Rogers-Ramanujan identities as evaluations of Ramanujan's function defined by

$$A_q(z) = \sum_{n=0}^{\infty} \frac{q^n}{(q;q)_n}(-z)^n.$$

Namely, (1.1) evaluates $A_q(-1)$, (1.2) evaluates $A_q(-q)$ .

Another remarkable identity, which is even simpler, is the $q$-binomial identity [6, (II.3)]

$$\sum_{n=0}^{\infty} \frac{(a;q)_n}{(q;q)_n} z^n = \frac{(az;q)_\infty}{(z;q)_\infty}. \tag{1.3}$$

In recent work by Ismail and zhang [8], the following function was considered

$$A_q^{(\alpha)}(a;t) = \sum_{n=0}^{\infty} \frac{(a;q)_n \, q^{\alpha n^2} t^n}{(q;q)_n}. \tag{1.4}$$

Clearly, $A_q^{(\alpha)}(a;t)$ specializes to $A_q(z)$ with $a = 0, \alpha = 1$ and $t = -z$. Furthermore, it specializes to the left hand side of (1.3) when $\alpha = 0$. Naturally, one wouldn't expect a simple closed form identity for such a general function, nonetheless we present in section 2 below some generalizations of identities on $A_q^{(\alpha)}$ relating different values of the parameter $\alpha$. We also address similar generalizations to a bilateral analogue of (1.4) related to the Ramanujan ${}_1\psi_1$ sums [6, (II.29)]

$$\sum_{-\infty}^{\infty} \frac{(a;q)_n}{(b;q)_n} z^n = \frac{(q, b/a, az, q/az; q)_\infty}{(b, q/a, z, b/az; q)_\infty}, \quad \left|\frac{b}{a}\right| < |z| < 1. \tag{1.5}$$

In section 3, we consider the series

$$F(a, c; z) := \sum_{k=0}^{\infty} \frac{(a;q)_k (-1)^k q^{k(k-1)/2} z^k}{(q, c; q)_k},$$

which can be obtained as a limit of basic hypergeometric series. We then utilize hypergeometric transformations to obtain some infinite and terminating series identites.

# 2 Unilateral and Bilateral analogues of Rogers-Ramanujan Identities

We start by recalling the following results which were proved in [8].

**Lemma 2.1.** *[8, Lemma 4.1] For nonnegative integer $j,k,\ell,m,n$ and $\rho=e^{2\pi i/3}$ we have*

$$\sum_{k=0}^{n}\frac{(a;q)_k\,(a;q)_{n-k}\,(-1)^k}{(q;q)_k\,(q;q)_{n-k}}=\begin{cases}0 & n=2m+1\\ \frac{\left(a^2;q^2\right)_m}{\left(q^2;q^2\right)_m} & n=2m\end{cases}, \tag{2.1}$$

*and*

$$\sum_{\substack{j+k+\ell=n\\ j,k,\ell\geq 0}}\frac{(a;q)_j\,(a;q)_k\,(a;q)_\ell}{(q;q)_j\,(q;q)_k\,(q;q)_\ell}\rho^{k+2\ell}=\begin{cases}0 & 3\nmid n\\ \frac{\left(a^3;q^3\right)_m}{\left(q^3;q^3\right)_m} & n=3m\end{cases}. \tag{2.2}$$

*For $j,k,m,\ell,n\in F$, we have*

$$\sum_{j+k=n}\frac{(a;q)_j\,(a;q)_k\,(-1)^k}{(b;q)_j\,(b;q)_k}=\begin{cases}0 & n=2m+1\\ \frac{(q,b/a,-b,-q/a;q)_\infty}{(-q,-b/a,b,q/a;q)_\infty}\frac{\left(a^2;q^2\right)_m}{\left(b^2;q^2\right)_m} & n=2m\end{cases} \tag{2.3}$$

*and*

$$\sum_{j+k+\ell=n}\frac{(a;q)_j\,(a;q)_k\,(a;q)_\ell\,\rho^{k+2\ell}}{(b;q)_j\,(b;q)_k\,(b;q)_\ell}=0 \tag{2.4}$$

*for $3\nmid n$,*

$$\begin{aligned}&\sum_{j+k+\ell=3m}\frac{(a;q)_j\,(a;q)_k\,(a;q)_\ell\,\rho^{k+2\ell}}{(b;q)_j\,(b;q)_k\,(b;q)_\ell}\\ =\;&\frac{(q,b/a;q)_\infty^3}{(b,q/a;q)_\infty^3}\frac{(b^3,q^3a^{-3};q^3)_\infty}{(q^3,b^3a^{-3};q^3)}\frac{(a^3;q^3)_m}{(b^3;q^3)_m}.\end{aligned} \tag{2.5}$$

We start by proving an extension of these results to general primitive roots of unity. In order to make our notation compact; we define for $r\geq 2$ and $n\geq 0$ sets

$$C_r(n):=\left\{(k_1,\dots,k_r):k_i\in F\text{ and }\sum_{i=1}^{r}k_i=n\right\}. \tag{2.6}$$

Also, let $C_r^+(n)$ be the subset of $C_r(n)$ whose entries are all nonnegative. Furthermore, we let $\zeta_r$ denote the primitive $r$th root of unity $e^{2\pi i/r}$.

**Lemma 2.2.** *The following identities hold for $r\geq 2$*

$$\sum_{(k_1,\dots,k_r)\in C_r^+(n)}\frac{(a;q)_{k_1}\,(a;q)_{k_2}\cdots(a;q)_{k_r}}{(q;q)_{k_1}\,(q;q)_{k_2}\cdots(q;q)_{k_r}}\zeta_r^{\sum_{i=1}^r ik_i}=\begin{cases}0 & r\nmid n,\\ \frac{(a^r;q^r)_m}{(q^r;q^r)_m} & n=rm,\end{cases} \tag{2.7}$$

$$\sum_{(k_1,\dots,k_r)\in C_r(n)}\frac{(a;q)_{k_1}\cdots(a;q)_{k_r}}{(b;q)_{k_1}\cdots(b;q)_{k_r}}\left(\zeta_r\right)^{\sum_{i=1}^r ik_i}=\begin{cases}0 & r\nmid n,\\ \frac{(q,b/a;q)_\infty^r}{(b,q/a;q)_\infty^r}\frac{\left(b^r,q^ra^{-r};q^r\right)_\infty}{\left(q^r,b^ra^{-r};q^r\right)}\frac{(a^r;q^r)_m}{(b^r;q^r)_m} & n=rm,\end{cases} \tag{2.8}$$

*Proof.* We start by noting that, for $|t|<1$ we have

$$\frac{(at;q)_\infty}{(t;q)_\infty}\frac{(a\zeta_r t;q)_\infty}{(\zeta_r t;q)_\infty}\cdots\frac{(a\zeta_r^{r-1}t;q)_\infty}{(\zeta_r^{r-1}t;q)_\infty}=\frac{(a^r t^r;q^r)}{(t^r;q^r)},\quad |t|<1.$$

Employing (1.3) we see that

$$\prod_{i=0}^{r-1}\left(\sum_{k_i=0}^{\infty}\frac{(a;q)_{k_i}}{(q;q)_{k_i}}(\zeta_r^i x)^{k_i}\right)=\frac{(a^r x^r;q^r)}{(x^r;q^r)}=\sum_{m=0}^{\infty}\frac{(a^r;q^r)_m}{(q^r;q^r)_m}x^{rm},$$

and (2.7) follows by comparing the coefficients of $x^n$ in (2.9).

The proof of (2.8) is similar. We start by noting that for $|ba^{-1}|<|x|<1$, we have

$$\prod_{i=0}^{r-1}\frac{(q,b/a,a\zeta_r^i z,q\zeta_r^{-i}/az;q)_\infty}{(b,q/a,\zeta_r^i z,b\zeta_r^{-i}/az;q)_\infty}=\frac{(q,b/a;q)_\infty^r}{(b,q/a;q)_\infty^r}\frac{(a^r z^r,q^r a^{-r}z^{-r};q^r)_\infty}{(z^r,b^r a^{-r}z^{-r};q^r)_\infty}$$

applying the Ramanujan ${}_1\psi_1$ sum (1.5) to that identity establishes (2.8). □

Lemma (2.2) enables us to prove the following result generalizing Theorem 4.2 of [8]

**Theorem 2.3.** *For $\alpha\geq 0$ and integer $r\geq 2$ we have*

$$A_{q^r}^{(r\alpha)}(a^r;t^r)=\sum_{k_1,\ldots,k_{r-1}=0}^{\infty}\frac{(a;q)_{k_1}\cdots(a;q)_{k_{r-1}}\zeta_r^{\sum_{i=1}^{r-1}ik_i}q^{\alpha\left(\sum_{i=1}^{r-1}k_i\right)^2}t^{\sum_{i=1}^{r-1}k_i}}{(q;q)_{k_1}\cdots(q;q)_{k_{r-1}}}A_q^{(\alpha)}\left(a;q^{2\alpha(\sum_{i=1}^{r-1}k_i)}t\right). \tag{2.9}$$

$$A_{q^r}^{(r\alpha)}(a^r;t^r)=\sum_{k_2,\ldots,k_r=0}^{\infty}\frac{(a;q)_{k_2}\cdots(a;q)_{k_r}\zeta_r^{\sum_{i=2}^{r-1}ik_i}q^{\alpha\left(\sum_{i=2}^{r}k_i\right)^2}t^{\sum_{i=2}^{r}k_i}}{(q;q)_{k_2}\cdots(q;q)_{k_r}}A_q^{(\alpha)}\left(a;\zeta_r q^{2\alpha(\sum_{i=2}^{r}k_i)}t\right). \tag{2.10}$$

*Proof.* The proof of (2.9)follows from the following series identities

$$\begin{aligned}A_{q^r}^{(r\alpha)}(a^r;t^r)&=\sum_{m=0}^{\infty}\frac{(a^r;q^r)_m}{(q^r;q^r)_m}q^{\alpha r^2m^2}t^{rm}\\&=\sum_{n=0}^{\infty}\sum_{(k_1,\ldots,k_r)\in C_r^+(n)}\frac{(a;q)_{k_1}\cdots(a;q)_{k_r}}{(q;q)_{k_1}\cdots(q;q)_{k_r}}\zeta_r^{\sum_{i=1}^{r}ik_i}t^n q^{\alpha n^2}\\&=\sum_{k_1,\ldots,k_{r-1}=0}^{\infty}\frac{(a;q)_{k_1}\cdots(a;q)_{k_{r-1}}\zeta_r^{\sum_{i=1}^{r-1}ik_i}q^{\alpha\left(\sum_{i=1}^{r-1}k_i\right)^2}t^{\sum_{k=1}^{r-1}k_i}}{(q;q)_{k_1}\cdots(q;q)_{k_{r-1}}}A_q^{(\alpha)}\left(a;tq^{2\alpha\left(\sum_{i=1}^{r-1}k_i\right)}\right).\end{aligned} \tag{2.11}$$

The proof of (2.10) is almost identical, except that in the last step the inner sum is performed over $k_1$ rather than $k_r$. □

Following [8] we also consider the following generalization of the ${}_1\psi_1$ function. For $\alpha \geq 0$, define $B_q^{(\alpha)}$ by

$$B_q^{(\alpha)}(a,b;x) = \sum_{n=-\infty}^{\infty} \frac{(a;q)_n}{(b;q)_n} q^{\alpha n^2} x^n. \tag{2.12}$$

Note that $B_q^{(\alpha)}$ is a bilateral series analogue of $A_q^{(\alpha)}$ (with one additional parameter in fact as a result of the generality afforded by the ${}_1\psi_1$ formula). Again using Lemma (2.2) we obtain the following result.

**Theorem 2.4.** *We have*

$$\begin{aligned} B_{q^r}^{(r\alpha)}(a^r,b^r;x^r) &= \frac{(b,q/a;q)_\infty^r}{(q,b/a;q)_\infty^r} \frac{(q^r,b^r a^{-r};q^r)_\infty}{(b^r,q^r a^{-r};q^r)_\infty} \\ &\times \sum_{k_1,\ldots,k_{r-1}=-\infty}^{\infty} \frac{(a;q)_{k_1}\cdots(a;q)_{k_{r-1}}}{(b;q)_{k_1}\cdots(b;q)_{k_{r-1}}} \zeta_r^{\sum_{i=1}^{r-1} ik_i} q^{\alpha\left(\sum_{i=1}^{r-1} k_i\right)^2} x^{\sum_{k=1}^{r-1} k_i} B_q^{(\alpha)}\left(a,b;xq^{2\alpha\left(\sum_{i=1}^{r-1} k_i\right)}\right). \end{aligned} \tag{2.13}$$

*Proof.* The proof follows from the following series identities

$$\begin{aligned} &\frac{(q,b/a;q)_\infty^r}{(b,q/a;q)_\infty^r} \frac{(b^r,q^r a^{-r};q^r)_\infty}{(q^r,b^r a^{-r};q^r)_\infty} B_{q^r}^{(r\alpha)}(a^r,b^r;x^r) \\ &= \frac{(q,b/a;q)_\infty^r}{(b,q/a;q)_\infty^r} \frac{(b^r,q^r a^{-r};q^r)_\infty}{(q^r,b^r a^{-r};q^r)_\infty} \sum_{m=-\infty}^{\infty} \frac{(a^r;q^r)_m}{(b^r;q^r)_m} q^{\alpha r^2 m^2} x^{rm} \\ &= \sum_{n=-\infty}^{\infty} \sum_{(k_1,\ldots,k_r)\in C_r(n)} \frac{(a;q)_{k_1}\cdots(a;q)_{k_r}}{(b;q)_{k_1}\cdots(b;q)_{k_r}} \zeta_r^{\sum_{i=1}^{r} ik_i} x^n q^{\alpha n^2} \\ &= \sum_{k_1,\ldots,k_{r-1}=-\infty}^{\infty} \frac{(a;q)_{k_1}\cdots(a;q)_{k_{r-1}}}{(b;q)_{k_1}\cdots(b;q)_{k_{r-1}}} \zeta_r^{\sum_{i=1}^{r-1} ik_i} q^{\alpha\left(\sum_{i=1}^{r-1} k_i\right)^2} x^{\sum_{k=1}^{r-1} k_i} B_q^{(\alpha)}\left(a,b;xq^{2\alpha\left(\sum_{i=1}^{r-1} k_i\right)}\right). \end{aligned} \tag{2.14}$$

□

**Corollary 2.5.** *The following family of bilateral Rogers-Ramanujan type identities hold*

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \frac{q^{r^2n^2} x^{rn}}{1-a^r q^{rn}} &= \frac{(q^r;q^r)_\infty}{(q;q)_\infty^r} \frac{(a,q/a;q)_\infty^r}{(a^r,q^r a^{-r};q^r)_\infty} \\ &\times \sum_{k_1,\ldots,k_r=-\infty}^{\infty} \frac{\zeta_r^{\sum ik_i} q^{(\sum k_i)^2} x^{\sum k_i}}{(1-aq^{k_1})\cdots(1-aq^{k_r})}. \end{aligned} \tag{2.15}$$

*Proof.* Setting $\alpha = 1$ and $b = aq$ and using (2.13) we see that

$$\begin{aligned} B_{q^r}^{(r)}(a^r, a^r q^r; x^r) &= \frac{(q,q;q)_\infty^r}{(qa, q/a; q)_\infty^r} \frac{(q^r a^r, q^r a^{-r}; q^r)_\infty}{(q^r, q^r; q^r)_\infty} (1-a^r) \sum_{m=0}^{\infty} \frac{q^{\alpha r^2 m^2} x^{rm}}{(1-a^r q^m)} \\ &= (1-a)^r \sum_{k_1,\ldots,k_r=-\infty}^{\infty} \frac{\zeta_r^{\sum_{i=1}^r i k_i} x^{\sum k_i} q^{\alpha(\sum k_i)^2}}{(1-aq^{k_1}) \cdots (1-aq^{k_r})}, \end{aligned} \tag{2.16}$$

and the result follows on simplification. □

# 3 Heine Transforms and related identities

We consider the basic hypergeometric function defined by

$${}_2\phi_1 \left( \begin{matrix} a, b \\ c \end{matrix} \middle| q, z \right) := \sum_{k=0}^{\infty} \frac{(a;q)_k (b;q)_k}{(c;q)_k (q;q)_k} z^k. \tag{3.1}$$

We shall study the function

$$F(a, c; z) := \sum_{k=0}^{\infty} \frac{(a;q)_k (-1)^k q^{k(k-1)/2} z^k}{(q, c; q)_k} = \lim_{b\to\infty} {}_2\phi_1 \left( \begin{matrix} a, b \\ c \end{matrix} \middle| q, \frac{z}{b} \right). \tag{3.2}$$

where we have utilized the limit

$$\lim_{b\to\infty} \frac{(b;q)_k}{b^k} = (-1)^k q^{k(k-1)/2}.$$

We have the following alternative representation of $F(a; c; z)$

**Theorem 3.1.** *For $c, z \neq q^{-m}$ ($m$ nonnegative integer) we have*

$$F(a, c, z) = \frac{(z;q)_\infty}{(c;q)_\infty} \sum_{k=0}^{\infty} \frac{(az/c;q)_k (-1)^k q^{k(k-1)/2} c^k}{(q;q)_k (z;q)_k} \tag{3.3}$$

*Proof.* Recall the Heine transformation [6, (III.2)]

$${}_2\phi_1 \left( \begin{matrix} A, B \\ C \end{matrix} \middle| q, z \right) = \frac{(C/B, Bz; q)_\infty}{(C, z; q)_\infty} {}_2\phi_1 \left( \begin{matrix} ABz/C, B \\ Bz \end{matrix} \middle| q, \frac{C}{B} \right). \tag{3.4}$$

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{(a;q)_k (-1)^k q^{k(k-1)/2} z^k}{(q; c; q)_k} &= \lim_{b\to\infty} {}_2\phi_1 \left( \begin{matrix} a, b \\ c \end{matrix} \middle| q, \frac{z}{b} \right) \\ &= \lim_{b\to\infty} \frac{(c/b;q)_\infty (z;q)_\infty}{(c;q)_\infty (z/b;q)_\infty} {}_2\phi_1 \left( \begin{matrix} az/c, b \\ z \end{matrix} \middle| q, \frac{c}{b} \right) \\ &= \frac{(z;q)_\infty}{(c;q)_\infty} \sum_{k=0}^{\infty} \frac{(az/c;q)_k (-1)^k q^{k(k-1)/2} c^k}{(q;q)_k (z;q)_k} \end{aligned}$$

For the transformation formula to be valid we need neither $c$ nor $z$ to be of the form $q^{-m}$. This proves (3.3). □

**Remark.** Note that for $c = z$, both sides of (3.3) are identical. Otherwise the result above provides a way of interchanging the roles of $c$ and $z$ in which both sums are very similar except that the right hand side acquires the factor $\frac{(z;q)_\infty}{(c;q)_\infty}$ and the expression $(a;q)_k$ is replaced by $(az/c;q)_k$.

In the following corollary, we examine a case in which the above Theorem leads to transforming infinite sums into finite ones.

**Corollary 3.2.** *Let $\alpha, \gamma$ be real numbers such that none of $\alpha+\gamma$ and $\gamma-n$ are negative integers for any nonnegative integer $n$ (in particular, $\gamma$ itself can not be a negative integer). Then the following identity holds*

$$\begin{aligned} &\sum_{k=0}^{\infty} \frac{(q^\alpha;q)_k(-1)^k q^{k(k-1)/2} q^{k(\gamma-n)}}{(q, q^{\alpha+\gamma};q)_k} \\ &= \frac{(q^{\gamma-n};q)_\infty}{(q^{\alpha+\gamma};q)_\infty} \sum_{k=0}^{n} \frac{(q^{-n};q)_k(-1)^k q^{k(k-1)/2} q^{k(\gamma+\alpha)}}{(q;q)_k(q^{\gamma-n};q)_k}, \end{aligned} \tag{3.5}$$

*Proof.* This follows directly from (3.3) by setting $a = q^\alpha$ and $c = q^\gamma$ to obtain

$$F(q^\alpha, q^{\alpha+\gamma}; q^{\gamma-n}) = \frac{(q^{\gamma-n};q)_\infty}{(q^{\alpha+\gamma};q)_\infty} \sum_{k=0}^{n} \frac{(q^{-n};q)_k(-1)^k q^{k(k-1)/2} q^{k(\gamma+\alpha)}}{(q;q)_k(q^{\gamma-n};q)_k}. \tag{3.6}$$

□

A well-known consequence of the $q$-binomial theorem (1.3) for $|x| < 1$ (see p. 490 of [4] for instance)

$$\frac{1}{(x;q)_n} = \sum_{k=0}^{\infty} \frac{(q;q)_{n+k-1}}{(q;q)_k(q;q)_{n-1}} x^k. \tag{3.7}$$

Combining (3.7) with our results above yields the following evaluation.

**Corollary 3.3.** *For $|x| < 1$ and integer $n \geq 0$ we have*

$$\sum_{k=0}^{\infty} \frac{(q;q)_{n+k-1}}{(q;q)_k(q;q)_{n-1}} x^k = \sum_{k=0}^{n} \frac{(q^{-n};q)_k(-1)^k q^{k(k-1)/2}}{(q;q)_k(xq^{-n};q)_k} x^k. \tag{3.8}$$

*Proof.* By setting $\alpha = 0$ in (3.5) the left hand side simplifies to 1 since $(1;q)_0 = 1$ and $(1;q)_k = 0$ for $k \geq 1$. We thus obtain

$$1 \;=\; \frac{(q^{\gamma-n};q)_\infty}{(q^\gamma;q)_\infty} \sum_{k=0}^{n} \frac{(q^{-n};q)_k(-1)^k q^{k(k-1)/2} q^{k\gamma}}{(q;q)_k(q^{\gamma-n};q)_k}. \tag{3.9}$$

Note that

$$\frac{(x;q)_\infty}{(xq^{-n};q)_\infty} = \frac{1}{(xq^{-n};q)_n}. \tag{3.10}$$

Thus, setting $x = q^\gamma$ in (3.9) and applying (3.7) we obtain (3.8). □

In fact, we can generalize the argument above to obtain the following family of formulas which exhibits a remarkable symmetry between two finite sums of different lengths

**Theorem 3.4.** *Let $m, n$ be nonnegative integers, and assume $x \neq q^l$ for any integer $l$, then the following holds*

$$\sum_{k=0}^{m} \frac{(q^{-m};q)_k (-1)^k q^{k(k-2n-1)/2} x^k}{(q, x/q^m;q)_k} = \frac{(xq^{-n};q)_\infty}{(xq^{-m};q)_\infty} \sum_{k=0}^{n} \frac{(q^{-n};q)_k (-1)^k q^{k(k-2m-1)/2} x^k}{(q, x/q^n;q)_k}.$$

*Proof.* This follows by setting $\alpha = -m$ in (3.5) and noting that $(q^{-m};q)_k = 0$ for $k \geq m+1$. It is well known that the expression $(y;q)_\infty$ with $|q| < 1$ converges, and since both sums are now finite there are no convergence concerns, except that we need to guarantee that $(x/q^m;q)_k$ (as well as $(x/q^n;q)_k$ and $(x/q^m;q)_\infty$ are nonzero), which leads to the restrictions placed on $x$. □

# References


[1] G. E. Andrews, *The Theory of Partitions*, Addison-Wesley, Reading, MA, 1976.

[2] G. E. Andrews, *q-series: Their development and application in analysis, number theory, combinatorics, physics, and computer algebra*, CBMS Regional Conference Series, Number 66, American Mathematical Society, Providence, RI, 1986.

[3] G. E. Andrews, A polynomial identity which implies the Rogers–Ramanujan identities, Scripta Math. **28** (1970), 297–305.

[4] G. E. Andrews, R. A. Askey, and R. Roy, *Special Functions*, Cambridge University Press, Cambridge, 1999.

[5] K. Garrett, M. E. H. Ismail, and D. Stanton, Variants of the Rogers–Ramanujan identities, Advances in Applied Math. **23** (1999), 274–299.

[6] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, second edition, Encyclopedia of Mathematics and Its Applications, volume 96 Cambridge University Press, Cambridge, 2004.

[7] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in one Variable*, paperback edition, Cambridge University Press, Cambridge, 2009.

[8] M. E. H. Ismail and R. zhang, $q$-Bessel functions and Rogers-Ramanujan type identities, to appear in Proc. Amer. Math. Soc.

[9] L. J. Slater, *Generalized Hypergeometric Series*, Cambridge University Press, Cambridge, 1964.

[10] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers* (Introduction by G. E. Andrews), Narosa, New Delhi, 1988.

Ahmad El-Guindy
Current address: Science Program, Texas A&M University in Qatar, Doha, Qatar
Permanent address: Department of Mathematics, Faculty of Science, Cairo University, Giza, Egypt 12613
email: a.elguindy@gmail.com

M. E. H. I, University of Central Florida, Orlando, Florida 32828,
and King Saud University, Riyadh, Saudi Arabia
email: ismail@math.ucf.edu